

Shift of $^2S_{1/2}$ hyperfine splittings due to blackbody radiation

Wayne M. Itano, L. L. Lewis, and D. J. Wineland

Frequency and Time Standards Group, Time and Frequency Division, National Bureau of Standards, Boulder, Colorado 80303

(Received 27 October 1981)

Frequency shifts of hyperfine splittings of $^2S_{1/2}$ states due to the blackbody radiation field are calculated. It is shown that they can be estimated from the dc hyperfine Stark shifts, which have previously been measured in the ground states of hydrogen and the alkali atoms. The shift at 300 K is large enough to be significant in primary Cs atomic beam frequency standards, and should be measurable. A simple method of calculating the hyperfine Stark shifts is described, which is based on the Bates-Damgaard method for determining radial matrix elements and the Fermi-Segrè formula for determining the contact hyperfine matrix elements. It is applied to Ba^+ and Hg^+ , for which no experimental data are yet available, and which are currently of interest for frequency standards.

The most accurate and stable atomic frequency standards are based on hyperfine transition frequencies in $^2S_{1/2}$ ground states, such as in ^{133}Cs , 1H , and ^{87}Rb . In this Communication, we estimate the temperature-dependent shift of $^2S_{1/2}$ hyperfine splittings due to the blackbody radiation field. We note that this effect is large enough to be observable in a Cs atomic beam apparatus. The shift of the Cs hyperfine splitting at $T = 300$ K from the unperturbed ($T = 0$ K) value causes a frequency offset which is significant for primary frequency standards.

According to the Planck radiation law,

$$E^2(\omega)d\omega = B^2(\omega)d\omega = \frac{8\alpha^3}{\pi} \frac{\omega^3 d\omega}{\exp(\omega/kT) - 1}, \quad (1)$$

where $E^2(\omega)d\omega [B^2(\omega)d\omega]$ is the squared amplitude of the blackbody electric (magnetic) field in a bandwidth $d\omega$ around ω . [Atomic units (a.u.) are used unless otherwise specified ($\hbar = m_e = e = 1$).] The mean-squared fields are

$$\begin{aligned} \langle E^2(t) \rangle &= \frac{1}{2} \int_0^\infty E^2(\omega) d\omega = \frac{4\pi^3}{15} \alpha^3 (kT)^4 \\ &= (8.319 \text{ V/cm})^2 [T(K)/300]^4 \end{aligned} \quad (2)$$

and

$$\begin{aligned} \langle B^2(t) \rangle &= \frac{1}{2} \int_0^\infty B^2(\omega) d\omega \\ &= (2.775 \times 10^{-6} \text{ T})^2 [T(K)/300]^4. \end{aligned} \quad (3)$$

Gallagher and Cooke¹ pointed out that these fields induce temperature-dependent shifts of transition frequencies in atoms and molecules through the ac Zeeman and Stark effects.^{2,3} They estimated the shifts in high- n Rydberg levels and some other systems. Blackbody frequency shifts have not yet been observed in any system.

The fractional blackbody ac Zeeman shift of the ground-state hyperfine splitting in H or Cs was estimated in Ref. 1 to be about 10^{-16} at $T = 300$ K. We have derived the following expression for this shift in any $^2S_{1/2}$ ground state, which is valid at zero dc magnetic field and at temperatures such that the peak of the blackbody spectrum is at a much higher frequency than the hyperfine frequency. We find

$$\begin{aligned} \frac{\delta\omega_{\text{hfs}}}{\omega_{\text{hfs}}} &\cong -\frac{\alpha^2}{24} (g_J - g_I)^2 \int_0^\infty \frac{B^2(\omega)}{\omega^2} d\omega \\ &= -(\pi/18) (g_J - g_I)^2 \alpha^5 (kT)^2 \\ &\cong -1.304 \times 10^{-17} [T(K)/300]^2, \end{aligned} \quad (4)$$

where g_J and g_I are the electronic and nuclear g factors, respectively. In the last line, we have assumed that $g_J = 2$ and that $|g_I/g_J| \ll 1$.

At laboratory temperatures, the blackbody ac Stark shift of the hyperfine splitting, which has previously been neglected, is generally larger than the ac Zeeman shift. The ac hyperfine Stark shift due to an electric field of frequency ω is approximately equal to the dc hyperfine Stark shift due to a static field with the same rms values, if $\omega \ll \omega_{\text{res}}$, where ω_{res} is the lowest allowed electric dipole transition frequency. The correction is of order $(\omega/\omega_{\text{res}})^2$ and will be considered in more detail below. For the ground states of any of the alkali atoms, $(\omega/\omega_{\text{res}})^2 \leq 3 \times 10^{-3}$, where ω is the frequency corresponding to the peak of the blackbody spectrum at 300 K. Therefore, at 300 K, the blackbody ac hyperfine shift is approximately equal to the shift caused by a dc field of 8.3 V/cm.

The dc hyperfine Stark shift was first observed in Cs by Haun and Zacharias.⁴ Later, it was observed in

H (see Ref. 5) and other alkali atoms.⁶ These experiments measured the Stark shift of the $(F = I + \frac{1}{2}, M_F = 0) \leftrightarrow (F = I - \frac{1}{2}, M_F = 0)$ transition. They can be considered to be measurements of the scalar hyperfine polarizabilities, which are independent of M_F and the orientation of the electric field, since the contributions from the tensor polarizabilities can be estimated and are less than the experimental uncertainties.⁷ Only the scalar polarizability contributes to the blackbody ac Stark shift, because of the isotropy of the blackbody radiation. The fractional ac Stark shift of the Cs hyperfine splitting can be estimated from the measured dc hyperfine polarizability (see Table I of Ref. 6) and Eq. (2) to be $-1.69(4) \times 10^{-14} [T(K)/300]^4$. This shift is large enough to affect the calibration of primary Cs frequency standards and therefore should be taken into account. For example, the fractional uncertainty of one primary Cs frequency standard (Cs 1 of the Physikalisch-Technische Bundesanstalt) is stated to be 6.5×10^{-15} , but has not been corrected for the blackbody shift.⁸

The theory of the dc hyperfine Stark shift of $^2S_{1/2}$ ground states is quite well developed, and the calculations are in good agreement with the experiments. The shift appears in the third order of perturbation theory, where the electric-field interaction is taken twice and the hyperfine interaction is taken once. For hydrogenic atoms and ions, an analytic solution has been obtained.^{9,10} Numerical calculations have been made for the neutral alkali atoms.¹¹⁻¹⁴ The shift in Li has been calculated without using perturbation theory by the spin-optimized self-consistent field method, with the electric field included in the Hamiltonian.¹⁵

We write the third-order perturbation expression for the scalar fractional dc hyperfine Stark shift of the $ns^2S_{1/2}$ state of an alkali-like atom or ion in the following form, which is independent of the spin and magnetic moment of the nucleus,

$$\delta\omega_{\text{hfs}}/\omega_{\text{hfs}} = -kE^2 = -(k_1 + k_2)E^2, \quad (5)$$

where

$$k_1 = \sum_{n''J} \frac{(n^2S_{1/2} || r || n''^2P_J)^2}{6[W(n''PJ) - W(nS)]^2} \quad (6a)$$

and

$$k_2 = - \sum_{n'',J,n' \neq n} \frac{(n^2S_{1/2} || r || n''^2P_J)}{3[W(n''PJ) - W(nS)]} \times \frac{(n^2S_{1/2} || r || n''^2P_J)\psi_{n''s}(0)}{[W(n'S) - W(nS)]\psi_{ns}(0)}. \quad (6b)$$

The reduced matrix elements of \vec{r} , the position operator of the valence electron, are defined with the conventions of Edmonds.¹⁶ We choose the phases of

the radial wave functions so that they are real. $W(n''PJ)$ and $W(nS)$ are the energies (not including the hyperfine interaction) of the n''^2P_J and the $ns^2S_{1/2}$ states, respectively, and $\psi_{ns}(0)$ is the value of $ns^2S_{1/2}$ wave function at the origin. The dc magnetic field is assumed to be so small that the Zeeman splitting is much less than the hyperfine splitting.

These formulas were derived by following the method of Feichtner *et al.*,¹³ in which the $ns^2S_{1/2}$ wave functions and energies are calculated to first order in the contact hyperfine interaction and the Stark shifts are then calculated by second-order perturbation theory. We ignore the spin-dipolar and quadrupole hyperfine interactions, since they contribute only to the tensor hyperfine polarizability. The terms k_1 and k_2 correspond, respectively, to the diagrams drawn in Figs. 1(a) and 1(b) of Lee *et al.*¹¹ and the "hfs" and "wave function" terms of Feichtner *et al.*¹³

We have developed a simple method of approximately evaluating Eqs. (6a) and (6b). We calculate the radial matrix elements using the Coulomb (Bates-Damgaard) approximation¹⁷ and the values of the s -state wave functions at the origin using the Fermi-Segré formula.¹⁸ We have used this method to calculate the scalar fractional ground-state hyperfine polarizabilities of Li, Na, K, Rb, and Cs and have obtained agreement with experiment to within 12% or better in all cases. The lowest three p states and the lowest five s states were included in the basis. In using the Bates-Damgaard method, the experimental term values were used to calculate the n^* values. For the p states, the centers of gravity of the two fine-structure levels were used for the term values. Table 3 of Ref. 17 was used with linear interpolation. In evaluating Eq. (6b), the phases of the wave functions must be kept consistent. If we define the s -state wave functions to be positive at the origin, then the sign of the radial integral involving the ns wave function, obtained from Table 3 of Ref. 17, must be reversed if n is even. This is because the Coulomb wave functions used by Bates and Damgaard, which are good approximations to the true wave functions outside the core, are positive for r greater than the last node [see Eq. (10) of Ref. 17] and the true ns wave functions have $n - 1$ nodes.

This method can be used for other atoms, for which no experimental data or calculations have yet been published, such as the singly ionized alkaline earths. We have carried out the calculations for the ground states of Hg^+ and Ba^+ , which are currently of interest for applications in stored-ion frequency standards.^{19,20} In atomic units, $k = 37.9$ for Hg^+ and $k = 902$ for Ba^+ . The conversion between atomic and laboratory units of E^2 is given by

$$E^2(\text{a.u.}) = 3.782 \times 10^{-20} E^2[(\text{V/cm})^2]. \quad (7)$$

At 300 K, the fractional blackbody ac hyperfine Stark Shifts are -9.9×10^{-17} and -2.4×10^{-15} for Hg^+ and Ba^+ , respectively. In rf trap experiments, the ac Stark shift due to the trapping fields may be larger.

The ground-state hyperfine shift due to an ac electric field of magnitude $E(t) = E(\omega) \cos \omega t$ can be obtained by the same method that was used to derive Eqs. (6a) and (6b), except that the formula for ac Stark shift is used. We assume that $\omega \ll \omega_{\text{res}}$, where ω_{res} is taken to be the frequency of the center of gravity of the fine-structure components of the first resonance line. If we assume that only the first excited p state makes a significant contribution, which is a good approximation for the alkali atoms, the shift is

$\delta\omega_{\text{hfs}}/\omega_{\text{hfs}}$

$$\begin{aligned} &= -\frac{1}{2} \{k_1[1 + 3(\omega/\omega_{\text{res}})^2] + k_2[1 + (\omega/\omega_{\text{res}})^2]\} E^2(\omega) \\ &= -k \langle E^2(t) \rangle [1 + (3k_1/k + k_2/k)(\omega/\omega_{\text{res}})^2] . \end{aligned} \quad (8)$$

The effect of the frequency distribution of the blackbody electric field is thus to increase $|\delta\omega_{\text{hfs}}/\omega_{\text{hfs}}|$ by the fractional amount

$$\epsilon = (3k_1/k + k_2/k) \langle \omega^2 \rangle / \omega_{\text{res}}^2 \quad (9)$$

relative to its value for a dc electric field of the same rms value. The mean-squared frequency of the

blackbody electric field $\langle \omega^2 \rangle$ is given by

$$\begin{aligned} \langle \omega^2 \rangle &= \int_0^\infty \omega^2 E^2(\omega) d\omega / \int_0^\infty E^2(\omega) d\omega \\ &= \frac{40\pi^2}{21} (kT)^2 . \end{aligned} \quad (10)$$

Among the alkali atoms, ϵ is the largest for Cs, which has the lowest value of ω_{res} . If we take k_1 and k_2 from Ref. 11, then for Cs,

$$\epsilon = 1.4 \times 10^{-2} [T(\text{K})/300]^2 , \quad (11)$$

which, at 300 K, is smaller than the experimental uncertainty in the dc hyperfine polarizability.

The blackbody shift could be observed in a Cs frequency standard which was modified so that the temperature of a tube surrounding the atoms in the resonance region could be varied. If the temperature were changed, for example, from 300 to 400 K, the fractional frequency shift would be 3.7×10^{-14} . If the frequency standard had the same frequency stability as NBS-6, the primary frequency standard of the United States,²¹ this shift could be determined to 30% or better in an averaging time of several hours.

This work was supported in part by the Air Force Office of Scientific Research and the Office of Naval Research.

¹T. F. Gallagher and W. E. Cooke, Phys. Rev. Lett. **42**, 835 (1979).

²M. Mizushima, Phys. Rev. **133**, A414 (1964).

³C. H. Townes and A. L. Schawlow, *Microwave Spectroscopy* (McGraw-Hill, New York, 1955), pp. 273–279.

⁴R. D. Haun and J. R. Zacharias, Phys. Rev. **107**, 107 (1957).

⁵E. N. Fortson, D. Kleppner, and N. F. Ramsey, Phys. Rev. Lett. **13**, 22 (1964); P. C. Gibbons and N. F. Ramsey, Phys. Rev. A **5**, 73 (1972); J. G. Stuart, D. J. Larson, and N. F. Ramsey, *ibid.* **22**, 2092 (1980).

⁶J. R. Mowat, Phys. Rev. A **5**, 1059 (1972).

⁷H. Gould, E. Lipworth, and M. C. Weisskopf, Phys. Rev. **188**, 24 (1969).

⁸G. Becker, IEEE Trans. Instrum. Meas. **29**, 297 (1980).

⁹C. Schwartz, Ann. Phys. (N.Y.) **6**, 156 (1959).

¹⁰P. G. H. Sandars, Proc. Phys. Soc. London **92**, 857 (1967).

¹¹T. Lee, T. P. Das, and R. M. Sternheimer, Phys. Rev. A **11**, 1784 (1975).

¹²N. L. Manakov, V. D. Ovsyannikov, and L. P. Rapoport, Opt. Spektrosk. **38**, 424 (1975) [Opt. Spectrosc. (USSR) **38**, 242 (1975)].

¹³J. D. Feichtner, M. E. Hoover, and M. Mizushima, Phys. Rev. **137**, A702 (1965).

¹⁴H. P. Kelly, R. L. Chase, G. R. Daum, and J. J. Chang,

Phys. Rev. A **8**, 2777 (1973).

¹⁵U. Kaldor, J. Phys. B **6**, 71 (1973).

¹⁶A. R. Edmonds, *Angular Momentum in Quantum Mechanics*, 2nd ed. (Princeton University, Princeton, N.J., 1974).

¹⁷D. R. Bates and A. Damgaard, Philos. Trans. R. Soc. London Ser. A **242**, 101 (1949).

¹⁸H. Kopfermann, *Nuclear Moments* (Academic, New York, 1958).

¹⁹F. G. Major and G. Werth, Phys. Rev. Lett. **30**, 1155 (1973); M. D. McGuire, R. Petsch, and G. Werth, Phys. Rev. A **17**, 1999 (1978); M. Jardino, M. Desaintfusien, R. Barillet, J. Viennet, P. Petit, and C. Audoin, Appl. Phys. **24**, 107 (1981); M. D. McGuire, Bull. Am. Phys. Soc. **26**, 615 (1981); D. J. Wineland, W. M. Itano, J. C. Bergquist, and F. L. Walls, in *Proceedings of the 35th Annual Symposium on Frequency Control* (U.S. Army Electronics Command, Fort Monmouth, N.J., 1981).

²⁰R. Blatt and G. Werth, Z. Phys. A **299**, 93 (1981); F. L. Walls, D. J. Wineland, and R. E. Drullinger, in *Proceedings of the 32nd Annual Symposium on Frequency Control* (U.S. Army Electronics Command, Fort Monmouth, N.J., 1978).

²¹D. J. Wineland, D. W. Allan, D. J. Glaze, H. W. Hellwig, and S. Jarvis, IEEE Trans. Instrum. Meas. **25**, 453 (1976).